



SIVANTO[®]

prime



Growing in unison

Product description:



- // Flupyradifurone is the active ingredient for **SIVANTO[®] prime** and was derived from an Asian plant *Stemona japonica*.
- // The exciting insecticidal properties come from Stemofoline a derivative of *Stemona japonica*.
- // **SIVANTO[®] prime** is a unique new chemistry that specifically targets major sucking pests.
- // **SIVANTO[®] prime** belongs to a new chemical class of Butenolides.

Proven safety:

- // **SIVANTO[®] prime** has been tested with highly bee attractive crops.
- // **SIVANTO[®] prime** has a low intrinsic toxicity to adult and immature honey bees.
- // **SIVANTO[®] prime** confers low risk to honey bees even when applied to flowering crops during bloom.
- // **SIVANTO[®] prime** is highly selective, safe to crops and enhances quality of harvest thus ensuring greater safety in agriculture.



Stemofoline




SIVANTO[®] prime benefits:

- // **SIVANTO[®] prime** provides convenience and flexibility as a foliar or drench application.
- // **SIVANTO[®] prime** causes quick feeding cessation and this provides effective virus vector control.
- // **SIVANTO[®] prime** has an excellent safety and environmental eco-toxicological profile.
- // **SIVANTO[®] prime** is a flexible systemic insecticide that controls key adult and immature sucking pests.
- // **SIVANTO[®] prime** is selective to most beneficials and a perfect fit for most integrated pest management (IPM) programs.

FIGURE 1


SIVANTO® prime 200 SL - Proven aphid control



Fast feeding cessation

- // **SIVANTO® prime** has excellent speed of action.
- // **SIVANTO® prime** causes quick feeding cessation.
- // **SIVANTO® prime** minimizes the probing by target pest and provides effective virus vector control.
- // **SIVANTO® prime** is selective for most beneficial insect pests.

Sivanto® prime caused an immediate feeding cessation effect and prevented both xylem and phloem feeding in Asian Citrus Psyllid (ACP) adults. ACP adults acquire and transmit bacteria that cause citrus greening disease when feeding on stems and leaves of citrus trees.



Application:

CROP	APPLICATION RATE PER HECTARE	REMARKS
TOBACCO Tobacco aphid <i>(Myzus P. Nicotianae)</i>	200 ml / 100 litres water	Scout for aphids once every week starting at 4 weeks after planting until topping. Apply Sivanto® prime as soon as aphids are noticed.

PLEASE NOTE: Read the Sivanto® prime label for more instructions.

SIVANTO® prime Reg. No. L10776 (Act No. 36 of 1947). **SIVANTO® prime** contains Flupyradifurone (Butenolide) (Caution). **Sivanto® prime** is registered trademark of Bayer AG, Germany. Use strictly according to instructions on label.

| @Bayer4Crops

Facebook: Bayer Crop Science Division Southern Africa **Twitter:** @bayer4cropssa
Bayer (Pty) Ltd. Reg. No. 1968/011192/07
 27 Wrench Road, Isando, 1601. PO Box 143, Isando, 1600, **Tel:** +27 11 921 5002

Bayer Zimbabwe: Farai Munyanyi, Territory Manager
Tel: +263 772 131228 **E-mail:** farai.munyanyi@bayer.com

www.cropscience.bayer.co.za
www.bayer.co.za

